

(가)

2006. 4. .

	()	가
	10, 20, 40 (, : pfizer Ireland Pharmaceuticals)	218
()	1 (10, 20, 40mg)	() 10.85, 21.69, 43.38 mg
	10mg, 20mg	
	1) : _____ , _____ (55 , , _____ HDL - _____ 가)가 가. _____	
	2) 2 _____ , _____ _____ (_____ , _____ , _____) _____)가 : 가. _____	
	3) _____ () (Type a b) , LDL - , , -B	
	4) _____ (Type IV)	
	5) _____ (Type III)	
	6) _____ 가 _____ 가 10 - 17 (_____ _____) , LDL - , -B 가. LDL - 190 mg/dL (190 mg/dL) . LDL - 160 mg/dL (160 mg/dL) , 가 가	

	<p style="text-align: center;">40mg</p> <p>1) _____ : _____ , _____ HDL- (55 _____ , _____)가 _____ 가. _____</p> <p>2) _____ 2 _____ , _____ _____ (_____ , _____ , _____) _____)가 _____ : _____ 가. _____</p> <p>3) _____ (_____) (Type a b) _____ , LDL- _____ , _____ -B</p> <p>4) _____ (Type IV)</p> <p>5) _____ (Type III)</p>
	<p>< 가 > 10mg, 20mg</p> <p>_____ 1 1 10mg . LDL- _____ , _____ 4 _____ .</p> <p>1) 80mg _____ (_____) _____ 1 1 10mg _____ . _____ 2 _____ 4 _____ .</p> <p>2) 가 _____ (10 ~ 17) _____ 1 10mg _____ , _____ 1 20mg _____ . (_____) _____ 20mg _____ .</p> <p>4 _____ .</p>

3)

LDL-
가 .

4)

70

40mg

1 1 10mg . LDL-

4

1

80mg

1)

()

1 1 10mg

2

4

2)

1 80mg

3)

LDL-
가 .

4)

70

< 가 >

(1~30) ,

36

- 가 ,

	- HMG-CoA reductase ()
	- 1997.9.30. 가 - 2005.9.21. 가 . <i>Prevention of Cardiovascular Disease</i> 가 . In patients with type 2 diabetes, and without clinically evident coronary heart disease, but with multiple risk factors for coronary heart disease such as retinopathy, albuminuria, smoking, or hypertension, LIPITOR is indicated to: ·Reduce the risk of myocardial infarction ·Reduce the risk of stroke
	3 2 7 가 : · 가 , (: 가)
	< > 1. · :
	:

· ()

·

가. ·

< 10 () >

1) :
, (55 ,
, HDL- 가)가

가.

2) , (,
,)가 2 :

가.

3) () (Type a b)
, LDL- , -B

4) 가 (Type IV)

5) (Type III)

6) 가

가 10 - 17 () ,
LDL- , -B

가. LDL- 190 mg/dL (190 mg/dL)

. LDL- 160 mg/dL (160 mg/dL) ,

가 가

가

< 20 () >

1) :
, (55 ,
, HDL- 가)가

가.

2) , ,)가 2 : (,

가.

3) () (Type a b) , LDL- , -B

4) 가 (Type IV)

5) (Type III)

6) 가 () ,

가 10 - 17 () , LDL- -B

가. LDL- 190 mg/dL (190 mg/dL)

. LDL- 160 mg/dL (160 mg/dL) ,

가 가

가

< 40 () >

1) : , (55 , HDL- 가)가

가.

2) , ,)가 2 : (,

가.

3) () (Type a b) , LDL- , -B

4) 가 (Type IV)

5) (Type III).