

< .

>

: . (2003-17 ,
2003.4.14.) 3 2 7 가 (.)

	가	
	()	()
	10, 20, 40mg	10, 20, 40mg
	1 () 10.85, 21.69, 43.38 mg (10, 20, 40mg)	1 () 10.85, 21.69, 43.38 mg (10, 20, 40mg)
1)	10mg, 20mg : _____ : _____ _____, _____, HDL- (55 _____, _____, _____)가 _____) 가. _____	1) _____ : _____ _____, _____, HDL- (55 _____, _____, _____)가 _____) 가. _____
2)	(Type a b) () , LDL- -B ,	2) _____ , _____ _____, _____, _____)가 _____ : 가. _____
3)	(Type III)	3) _____ () (Type a b) () , LDL- -B ,
4)	가 가 10 - 17 (_____ , LDL-) -B	4) _____ (Type IV)
	가. LDL- 190 mg/dL (190 mg/dL) LDL- 160 mg/dL (160 mg/dL) , 가 가	5) _____ (Type III)
		6) 가 가 10 - 17 (_____ , LDL-) -B 가. LDL- 190 mg/dL (190 mg/dL) LDL- 160 mg/dL (160 mg/dL) , 가 가

	가	
	()	()
	10, 20, 40mg	10, 20, 40mg
	<p>1) 40mg : _____ , _____ (55 _____ , _____ HDL- _____ 가)가 가. _____ .</p> <p>2) _____ (Type a b) () _____ , LDL- _____ , _____ -B</p> <p>3) _____ (Type III)</p>	<p>1) 40mg : _____ , _____ (55 _____ , _____ HDL- _____ 가)가 가. _____ .</p> <p>2) 2 _____ , _____ _____ (_____ , _____) _____)가 : 가. _____ .</p> <p>3) _____ (Type a b) () _____ , LDL- _____ , _____ -B</p> <p>4) _____ (Type IV)</p> <p>5) _____ (Type III)</p>

1.

2.

- 1) A double-blind, placebo-controlled study of atorvastatin as prevention of CHD in high risk patients with non-insulin-dependent diabetes mellitus
Collaborative atorvastatin diabetes study (CARDS)
- 2) A 3- to 5 year double-blind, randomized, placebo-controlled study of atorvastatin as primary prevention of cardiovascular disease in high-risk with (type) noninsulin-dependent diabetes mellitus and micro- or macroalbuminuria (Protocol 981-119)
- 3) A 4-week, double-blind, placebo-controlled, multicenter study of once daily atorvastatin (CI-981) in patients with elevated triglycerides (protocol 981-38)

- 4) A multicenter, open-label, randomized, 12-week study comparing the efficacy and safety of atorvastatin (CI-981) with that of immediate release niacin in patients with elevated triglycerides (protocol 981-42)
- 5) A 24-week, randomized, open-label, parallel-arm, multicenter study to compare the efficacy and safety of atorvastatin versus fenofibrate in patients with elevated triglycerides (protocol 981-55)

3.

- 1) : Pfizer Ireland Pharmaceuticals "Lipitor 10mg, 20mg, 40mg, 80mg"