전문의약품
분류번호 : 239

디세텔®정 50

정신적인 불안정, 무절제한 식생활, 하제의 장기복용, 장관의 염증 등은 장관의 정상적인 운동을 파괴하며, 장관의 운동이 정상적이지 못하면 배변장애(변비, 설사, 교대성 변비 · 설사), 복통, 복부팽만감 등의 자각증상을 호소하게 됩니다.

일양디세텔정은 SOLVAY PHARMA사에서 개발한 Pinaverium Bromide를 주성분으로 하는 대장기능정상화제로서 비정상적인 장관운동 및 장관의 과민성을 정상화시켜 만성적인 장기능 장애를 개선시킵니다.

성 상
주황색 원형필름코팅정제

성분 · 함량
1정 중
브롬화 피나베리움(식약청고시 제98-127호) 50mg

약 리 작 용
1. 디세텔정은 First GI Selective Ca^{++} channel Antagonist로서 장관의 Hypermotility 및 Hypersensitivity를 정상화시킵니다.
2. 디세텔정은 통증, 복부팽만감, 설사, 변비, 설사와 변비의 교대 등 모든 IBS 증상 및 subtype 에 유효합니다.
3. 디세텔정은 장관의 L-type Ca^{++} channel에만 선택적으로 작용하고, 전신흡수율이 낮아 심혈관계 등의 부작용이 없습니다.
4. 디세텔정은 항콜린성 작용이 없으므로 녹내장, 전립선 환자 등의 대장 기능 이상에도 사용할 수 있습니다.

효능 · 효과
1. 주효능 효과
과민성대장증후군(복통, 복부팽만감, 복부경련, 변비, 설사, 변비와 설사의 반복, 구역, 구토), 담관운동부전.

용법 · 용량
성인 : 브롬화 피나베리움으로서 1회 50mg 1일 3회 식사중에 충분한 물과 함께 투여하십시오.
연령, 증상에 따라 적절히 증감하십시오.

사 용 상 의 주 의 사 항
1. 다음 환자에는 투여하지 마십시오.
임부 및 수유부
2. 부작용
1)소화기계 : 드물게 경미한 소화기장애가 나타날 수 있습니다.
2)과량투여 : 설사와 복부팽만감이 나타날 수 있습니다.
3. 상호작용
MAO저해제와의 병용투여는 피하는 것이 바람직합니다.

포 장 단 위
15, 500정

저 장 방 법
밀폐용기

※ 본 제품은 엄격한 품질관리를 필한 제품입니다.
만약 구입시 사용기한이 경과되었거나 변질 · 변패 또는 오손된 제품이 발견될 경우에는 구입하신 약국 또는 지점이나 사무소를 통하여 교환하여 드리며, 재정경제부가 고시한 "소비자피해보상규정"에 의거, 정당한 소비자피해에 대해 보상해 드립니다.

P0402-10

제조판매원
일양약품주식회사
경기도 용인시 기흥읍 하갈리 182-4
소비자상담실 : 080-021-1010(수신자요금부담전화)
학술상담실 : 031-270-6158 홈페이지 www.ilyang.co.kr

원료기술공급원
SOLVAY
SOLVAY PHARMACEUTICALS

dicetel® tabs. 50

■ Form and Presentation
Orange-colored, circular-shaped, film-coated tablet

■ Composition
Pinaverium Br 50mg

■ Indication
-Symptomatic treatment of pain, bowel disturbances, abdominal discomfort, constipation and diarrhea in related to irritable bowel syndrome (IBS)
-Symptomatic treatment of pain in related to functional disorders of the biliary tract

■ Dosage and Administration
The usual adult dosage is three tablets a day (one tablet 3 times a day).
It is recommended that the tablet be taken with a glass of water during meals or snacks.

■ Use in pregnancy
Reproductive studies performed in animals have not revealed the presence of teratogenic effects. However, the safety of DICETEL during pregnancy has not been established. Consequently, in the pregnant patient, this drug should only be administrated if, the judgment of the physician, its use is essential to the welfare of the patient.

■ Use during lactation
There have been no controlled studies in nursing women, therefore, the drug should be avoided during lactation

■ Adverse reactions
1) Rare cases of mild GI disorders
2) Symptoms of overdosage : In man, apart from diarrhea and/or flatulence, pinaverium bromide induced no undesirable effects in daily dosages of up to 1,200mg

■ Drug interactions
Avoid prescribing with MAO inhibitors

• Countries where pinaverium bromide is available

Algeria	Gabon	Peru
Argentina	Greece	Philippines
Austria	Guatemala	Poland
Bahrein	Guinea	Portugal
Belgium	Honduras	Saudi Arabia
Brazil	Hong Kong	Senegal
Bulgaria	Hungary	Slovakia
Burkina Faso	India	South Korea
Cameroon	Italy	Spain
Canada	Kuwait	Sudan
Central African Republic	Lebanon	Switzerland
Chile	Libya	Taiwan
China	Luxembourg	Thailand
Columbia	Madagascar	Togo
Congo	Mauritania	Tunisia
Cyprus	Mexico	Turkey
Czech Republic	Morocco	United Arab Emirates
Ecuador	Netherlands	Uruguay
Egypt	Oman	Vietnam
El Salvador	Panama	Panama
France	Paraguay	Paraguay